

Infinite-dimensional α -divergence minimisation for Variational Inference

Kamélia Daudel

University of Oxford
kamelia.daudel@stats.ox.ac.uk

OxCML seminar
19/11/2021

Joint work with Randal Douc and François Portier

Outline

- ① Introduction
- ② Infinite-dimensional α -divergence minimisation
- ③ Numerical experiments
- ④ Conclusion

Outline

- 1 Introduction
- 2 Infinite-dimensional α -divergence minimisation
- 3 Numerical experiments
- 4 Conclusion

Bayesian statistics

- Compute / sample from the **posterior density** of the latent variables y given the data \mathcal{D}

$$p(y|\mathcal{D}) = \frac{p(\mathcal{D}, y)}{p(\mathcal{D})}.$$

- Problem : for many important models, we can only evaluate $p(y|\mathcal{D})$ up to the constant $p(\mathcal{D})$.

Bayesian statistics

- Compute / sample from the **posterior density** of the latent variables y given the data \mathcal{D}

$$p(y|\mathcal{D}) = \frac{p(\mathcal{D}, y)}{p(\mathcal{D})}.$$

- Problem : for many important models, we can only evaluate $p(y|\mathcal{D})$ **up to the constant** $p(\mathcal{D})$.

Variational Inference in a nutshell

→ Variational Inference : inference is seen as an **optimisation problem**.

- ① Posit a *simpler* variational family \mathcal{Q} , where $q \in \mathcal{Q}$.
- ② Fit q to obtain the best approximation to the posterior density

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{|\mathcal{D}}),$$


where D is a measure of dissimilarity between the variational distribution \mathbb{Q} and the posterior distribution $\mathbb{P}_{|\mathcal{D}}$

→ Typically, D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : parametric family

(e.g. Mean-field)

$$\begin{cases} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k(\theta, y) : \theta \in \Theta\} \end{cases}$$



Variational Inference in a nutshell

→ Variational Inference : inference is seen as an **optimisation problem**.

- ① Posit a *simpler* variational family \mathcal{Q} , where $q \in \mathcal{Q}$.
- ② Fit q to obtain the best approximation to the posterior density

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{|\mathcal{D}}),$$


where D is a measure of dissimilarity between the variational distribution \mathbb{Q} and the posterior distribution $\mathbb{P}_{|\mathcal{D}}$

→ Typically, D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : parametric family

(e.g. Mean-field)

$$\begin{cases} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k(\theta, y) : \theta \in T\} \end{cases}$$



Variational Inference in a nutshell

→ Variational Inference : inference is seen as an **optimisation problem**.

- ① Posit a *simpler* variational family \mathcal{Q} , where $q \in \mathcal{Q}$.
- ② Fit q to obtain the best approximation to the posterior density

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{|\mathcal{D}}),$$


where D is a measure of dissimilarity between the variational distribution \mathbb{Q} and the posterior distribution $\mathbb{P}_{|\mathcal{D}}$

→ Typically, D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : parametric family

(e.g. Mean-field)

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k(\theta, y) : \theta \in T\} \end{array} \right.$$



Variational Inference in a nutshell

→ Variational Inference : inference is seen as an **optimisation problem**.

- ① Posit a *simpler* variational family \mathcal{Q} , where $q \in \mathcal{Q}$.
- ② Fit q to obtain the best approximation to the posterior density

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{|\mathcal{D}}),$$


where D is a measure of dissimilarity between the variational distribution \mathbb{Q} and the posterior distribution $\mathbb{P}_{|\mathcal{D}}$

→ Typically, D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : parametric family

(e.g. Mean-field)

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k(\theta, y) : \theta \in T\} \end{array} \right.$$



Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : Mean-field family

$$\begin{cases} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{cases}$$

- Can we select alternative/more general D ?
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?

Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)

divergence and \mathcal{Q} : Mean-field family

$$\begin{cases} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{cases}$$

- Can we select alternative/more general D ?
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?


Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)
divergence and \mathcal{Q} : Mean-field family

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in \Theta\} \end{array} \right.$$



- Can we select alternative/more general D ?
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?


Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)
divergence and \mathcal{Q} : Mean-field family

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{array} \right.$$



- Can we select alternative/more general D ?
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?


Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)
divergence and \mathcal{Q} : Mean-field family

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{array} \right.$$



- Can we select alternative/more general D ?
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?


Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)
divergence and \mathcal{Q} : Mean-field family

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{array} \right.$$



- Can we select alternative/more general D ?
→ D is the α -divergence
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?


Core question in Variational Inference

$$\inf_{q \in \mathcal{Q}} D(\mathbb{Q} || \mathbb{P}_{\mathcal{D}})$$

Question : How to choose D and \mathcal{Q} ?

→ D : exclusive Kullback-Leibler (KL)
divergence and \mathcal{Q} : Mean-field family

$$\left\{ \begin{array}{l} D_{KL}(\mathbb{Q} || \mathbb{P}) = \int_Y \log \left(\frac{q(y)}{p(y)} \right) q(y) \nu(dy) \\ \mathcal{Q} = \{q : y \mapsto k_1(\theta_1, y_1) k_2(\theta_2, y_2) : (\theta_1, \theta_2) \in T\} \end{array} \right.$$



- Can we select alternative/more general D ?
→ D is the α -divergence
- Can we design more expressive variational families \mathcal{Q} beyond traditional parametric families?
→ Put a prior on the variational parameter θ

Variational Inference with the α -divergence family

(Y, \mathcal{Y}, ν) : measured space, ν is a σ -finite measure on (Y, \mathcal{Y}) .

\mathbb{Q} and \mathbb{P} : $\mathbb{Q} \preceq \nu$, $\mathbb{P} \preceq \nu$ with $\frac{d\mathbb{Q}}{d\nu} = q$, $\frac{d\mathbb{P}}{d\nu} = p$.

α -divergence between \mathbb{Q} and \mathbb{P}

$$D_\alpha(\mathbb{Q} || \mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL),} \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL).} \end{cases}$$

Variational Inference with the α -divergence family

(Y, \mathcal{Y}, ν) : measured space, ν is a σ -finite measure on (Y, \mathcal{Y}) .

\mathbb{Q} and \mathbb{P} : $\mathbb{Q} \preceq \nu$, $\mathbb{P} \preceq \nu$ with $\frac{d\mathbb{Q}}{d\nu} = q$, $\frac{d\mathbb{P}}{d\nu} = p$.

α -divergence between \mathbb{Q} and \mathbb{P}


$$D_\alpha(\mathbb{Q} || \mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL)}, \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL)}. \end{cases}$$

- ① A **flexible** family of divergences...

Figure: In red, the Gaussian which minimises $D_\alpha(\mathbb{Q} || \mathbb{P})$ for a varying α



Adapted from V. Cevher's lecture notes (2008) <https://www.ece.rice.edu/~vc3/elec633/AlphaDivergence.pdf>

Variational Inference with the α -divergence family

α -divergence between \mathbb{Q} and \mathbb{P}

$$D_\alpha(\mathbb{Q}||\mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL),} \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL).} \end{cases}$$

- ① A **flexible** family of divergences...
- ② ...**suitable** for Variational Inference purposes...

$$\inf_{q \in \mathcal{Q}} D_\alpha(\mathbb{Q}||\mathbb{P}|_{\mathcal{D}}) = \inf_{q \in \mathcal{Q}} \Psi_\alpha(q; \mathbf{p})$$

$$\text{with } \Psi_\alpha(q; \mathbf{p}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy) \text{ and } \mathbf{p} = \mathbf{p}(\cdot, \mathcal{D})$$

Black-box alpha divergence minimization. J. Hernandez-Lobato et al. (2016). ICML

Rényi divergence variational inference. Y. Li and R. E Turner (2016). NeurIPS

Variational inference via χ -upper bound minimization A. Dieng et al. (2017). NeurIPS

Variational Inference with the α -divergence family

α -divergence between \mathbb{Q} and \mathbb{P}

$$D_\alpha(\mathbb{Q}||\mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL),} \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL).} \end{cases}$$

- ① A **flexible** family of divergences...
- ② ...**suitable** for Variational Inference purposes...

$$\inf_{q \in \mathcal{Q}} D_\alpha(\mathbb{Q}||\mathbb{P}|_{\mathcal{D}}) = \inf_{q \in \mathcal{Q}} \Psi_\alpha(q; \mathbf{p})$$

$$\text{with } \Psi_\alpha(q; \mathbf{p}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy) \text{ and } \mathbf{p} = \mathbf{p}(\cdot, \mathcal{D})$$

Black-box alpha divergence minimization. J. Hernandez-Lobato et al. (2016). ICML

Rényi divergence variational inference. Y. Li and R. E Turner (2016). NeurIPS

Variational inference via χ -upper bound minimization A. Dieng et al. (2017). NeurIPS

Variational Inference with the α -divergence family

α -divergence between \mathbb{Q} and \mathbb{P}

$$D_\alpha(\mathbb{Q}||\mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL),} \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL).} \end{cases}$$

- ① A **flexible** family of divergences...
- ② ...**suitable** for Variational Inference purposes...

$$\inf_{q \in \mathcal{Q}} D_\alpha(\mathbb{Q}||\mathbb{P}|_{\mathcal{D}}) = \inf_{q \in \mathcal{Q}} \Psi_\alpha(q; \mathbf{p})$$

with $\Psi_\alpha(q; p) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy)$ and $\mathbf{p} = p(\cdot, \mathcal{D})$

- ③ ...with good **convexity** properties : f_α is convex!

Variational Inference with the α -divergence family

α -divergence between \mathbb{Q} and \mathbb{P}

$$D_\alpha(\mathbb{Q}||\mathbb{P}) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy),$$

where

$$f_\alpha = \begin{cases} \frac{1}{\alpha(\alpha-1)} [u^\alpha - 1 - \alpha(u-1)], & \text{if } \alpha \in \mathbb{R} \setminus \{0, 1\}, \\ u \log(u) + 1 - u, & \text{if } \alpha = 1 \text{ (Exclusive KL),} \\ -\log(u) + u - 1, & \text{if } \alpha = 0 \text{ (Inclusive KL).} \end{cases}$$

- ① A **flexible** family of divergences...
- ② ...**suitable** for Variational Inference purposes...

$$\inf_{q \in \mathcal{Q}} D_\alpha(\mathbb{Q}||\mathbb{P}|_{\mathcal{D}}) = \inf_{q \in \mathcal{Q}} \Psi_\alpha(q; \mathbf{p})$$

with $\Psi_\alpha(q; p) = \int_Y f_\alpha \left(\frac{q(y)}{p(y)} \right) p(y) \nu(dy)$ and $\mathbf{p} = p(\cdot, \mathcal{D})$

- ③ ...with good **convexity** properties : f_α is convex!

Outline

- 1 Introduction
- 2 Infinite-dimensional α -divergence minimisation
- 3 Numerical experiments
- 4 Conclusion

Infinite-dimensional α -divergence minimisation

Infinite-dimensional gradient-based descent for alpha-divergence minimisation.

K. Daudel, R. Douc and F. Portier. Ann. Statist. 49 (4) 2250 - 2270, August 2021.

<https://doi.org/10.1214/20-AOS2035>.

Idea : Extend the traditional variational parametric family

$$\mathcal{Q} = \{y \mapsto k(\theta, y) : \theta \in \mathsf{T}\}$$

by putting a prior on the variational parameter θ

$$\mathcal{Q} = \left\{ q : y \mapsto \mu k(y) := \int_{\mathsf{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathsf{M} \right\}$$

and propose an update formula for μ that ensures a **systematic decrease** in the α -divergence at each step

$$\rightarrow \text{Finite Mixture Models} : \mu = \sum_{j=1}^J \lambda_j \delta_{\theta_j}$$

NB: The mapping $\mu \mapsto \Psi_\alpha(\mu k; p)$ is convex!

Infinite-dimensional α -divergence minimisation

Infinite-dimensional gradient-based descent for alpha-divergence minimisation.

K. Daudel, R. Douc and F. Portier. Ann. Statist. 49 (4) 2250 - 2270, August 2021.

<https://doi.org/10.1214/20-AOS2035>.

Idea : Extend the traditional variational parametric family

$$\mathcal{Q} = \{y \mapsto k(\theta, y) : \theta \in \mathsf{T}\}$$

by putting a prior on the variational parameter θ

$$\mathcal{Q} = \left\{ q : y \mapsto \mu k(y) := \int_{\mathsf{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathsf{M} \right\}$$

and propose an update formula for μ that ensures a **systematic decrease** in the α -divergence at each step

$$\rightarrow \text{Finite Mixture Models} : \mu = \sum_{j=1}^J \lambda_j \delta_{\theta_j}$$

NB: The mapping $\mu \mapsto \Psi_\alpha(\mu k; p)$ is convex!

Infinite-dimensional α -divergence minimisation

Infinite-dimensional gradient-based descent for alpha-divergence minimisation.

K. Daudel, R. Douc and F. Portier. Ann. Statist. 49 (4) 2250 - 2270, August 2021.

<https://doi.org/10.1214/20-AOS2035>.

Idea : Extend the traditional variational parametric family

$$\mathcal{Q} = \{y \mapsto k(\theta, y) : \theta \in \mathsf{T}\}$$

by putting a prior on the variational parameter θ

$$\mathcal{Q} = \left\{ q : y \mapsto \mu k(y) := \int_{\mathsf{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathsf{M} \right\}$$

and propose an update formula for μ that ensures a **systematic decrease** in the α -divergence at each step

$$\rightarrow \text{Finite Mixture Models} : \mu = \sum_{j=1}^J \lambda_j \delta_{\theta_j}$$

NB: The mapping $\mu \mapsto \Psi_\alpha(\mu k; p)$ is convex!

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k; p) \quad \text{with} \quad \Psi_\alpha(\mu k; p) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times \mathcal{Y}$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu, \alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu, \alpha} + \kappa))} \quad \text{with} \quad b_{\mu, \alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k; p) \quad \text{with} \quad \Psi_\alpha(\mu k; p) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times Y$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu, \alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu, \alpha} + \kappa))} \quad \text{with} \quad b_{\mu, \alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k; p) \quad \text{with} \quad \Psi_\alpha(\mu k; p) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times \mathcal{Y}$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu, \alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu, \alpha} + \kappa))} \quad \text{with} \quad b_{\mu, \alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k; p) \quad \text{with} \quad \Psi_\alpha(\mu k; p) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times \mathcal{Y}$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu, \alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu, \alpha} + \kappa))} \quad \text{with} \quad b_{\mu, \alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k) \quad \text{with} \quad \Psi_\alpha(\mu k) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times \mathcal{Y}$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu, \alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu, \alpha} + \kappa))} \quad \text{with} \quad b_{\mu, \alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

The (α, Γ) -descent algorithm

Optimisation problem

$$\inf_{\mu \in M} \Psi_\alpha(\mu k) \quad \text{with} \quad \Psi_\alpha(\mu k) := \int_Y f_\alpha \left(\frac{\mu k(y)}{p(y)} \right) p(y) \nu(dy)$$

- p is a **nonnegative measurable function** defined on (Y, \mathcal{Y})
- M is a subset of $M_1(T)$, the space of probability measures on T
- $K : (\theta, A) \mapsto \int_A k(\theta, y) \nu(dy)$ is a Markov transition kernel defined on $T \times \mathcal{Y}$ with density k

Algorithm

Let $\mu_1 \in M_1(T)$ be such that $\Psi_\alpha(\mu_1 k) < \infty$. The sequence of probability measures $(\mu_n)_{n \geq 1}$ is defined iteratively by

$$\mu_{n+1} = \mathcal{I}_\alpha(\mu_n), \quad n \geq 1$$

where for all $\mu \in M_1(T)$ and all $\theta \in T$,

$$\mathcal{I}_\alpha(\mu)(d\theta) = \frac{\mu(d\theta) \cdot \Gamma(b_{\mu,\alpha}(\theta) + \kappa)}{\mu(\Gamma(b_{\mu,\alpha} + \kappa))} \quad \text{with} \quad b_{\mu,\alpha}(\theta) = \int_Y k(\theta, y) f'_\alpha \left(\frac{\mu k(y)}{p(y)} \right) \nu(dy)$$

Conditions for a monotonic decrease

(A1) For all $(\theta, y) \in T \times Y$, $k(\theta, y) > 0$, $p(y) \geq 0$ and $\int_Y p(y)\nu(dy) < \infty$.

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

Theorem

Assume (A1) and (A2). Let $\mu \in M_1(T)$ be such that $\Psi_\alpha(\mu k) < \infty$ and $\mu(\Gamma(b_{\mu,\alpha} + \kappa)) < \infty$. Then,

- ① $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) \leq \Psi_\alpha(\mu k)$
- ② $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) = \Psi_\alpha(\mu k)$ if and only if $\mu = \mathcal{I}_\alpha(\mu)$

Conditions for a monotonic decrease

(A1) For all $(\theta, y) \in T \times Y$, $k(\theta, y) > 0$, $p(y) \geq 0$ and $\int_Y p(y)\nu(dy) < \infty$.

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

Theorem

Assume (A1) and (A2). Let $\mu \in M_1(T)$ be such that $\Psi_\alpha(\mu k) < \infty$ and $\mu(\Gamma(b_{\mu,\alpha} + \kappa)) < \infty$. Then,

- ① $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) \leq \Psi_\alpha(\mu k)$
- ② $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) = \Psi_\alpha(\mu k)$ if and only if $\mu = \mathcal{I}_\alpha(\mu)$

Conditions for a monotonic decrease

(A1) For all $(\theta, y) \in T \times Y$, $k(\theta, y) > 0$, $p(y) \geq 0$ and $\int_Y p(y)\nu(dy) < \infty$.

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

Theorem

Assume (A1) and (A2). Let $\mu \in M_1(T)$ be such that $\Psi_\alpha(\mu k) < \infty$ and $\mu(\Gamma(b_{\mu,\alpha} + \kappa)) < \infty$. Then,

- ① $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) \leq \Psi_\alpha(\mu k)$
- ② $\Psi_\alpha(\mathcal{I}_\alpha(\mu)k) = \Psi_\alpha(\mu k)$ if and only if $\mu = \mathcal{I}_\alpha(\mu)$

Examples satisfying (A2)

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

- Entropic Mirror Descent : $\eta \in (0, 1]$, $\kappa \in \mathbb{R}$ and $\alpha = 1$

$$\Gamma(v) = e^{-\eta v}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \exp \left[-\eta \int_Y k(\theta, y) \log \left(\frac{\mu_n k(y)}{p(y)} \right) \nu(\mathrm{d}y) \right]$$

→ NB : η corresponds to the learning rate

- Power descent : $\eta \in (0, 1]$, $(\alpha - 1)\kappa \geq 0$ and $\alpha \neq 1$

$$\Gamma(v) = [(\alpha - 1)v + 1]^{\frac{\eta}{1-\alpha}}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \left[\int_Y k(\theta, y) \left(\frac{\mu_n k(y)}{p(y)} \right)^{\alpha-1} \nu(\mathrm{d}y) + (\alpha - 1)\kappa \right]^{\frac{\eta}{1-\alpha}}$$

Examples satisfying (A2)

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

- Entropic Mirror Descent : $\eta \in (0, 1]$, $\kappa \in \mathbb{R}$ and $\alpha = 1$

$$\Gamma(v) = e^{-\eta v}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \exp \left[-\eta \int_Y k(\theta, y) \log \left(\frac{\mu_n k(y)}{p(y)} \right) \nu(\mathrm{d}y) \right]$$

→ NB : η corresponds to the learning rate

- Power descent : $\eta \in (0, 1]$, $(\alpha - 1)\kappa \geq 0$ and $\alpha \neq 1$

$$\Gamma(v) = [(\alpha - 1)v + 1]^{\frac{\eta}{1-\alpha}}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \left[\int_Y k(\theta, y) \left(\frac{\mu_n k(y)}{p(y)} \right)^{\alpha-1} \nu(\mathrm{d}y) + (\alpha - 1)\kappa \right]^{\frac{\eta}{1-\alpha}}$$

Examples satisfying (A2)

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

- Entropic Mirror Descent : $\eta \in (0, 1]$, $\kappa \in \mathbb{R}$ and $\alpha = 1$

$$\Gamma(v) = e^{-\eta v}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \exp \left[-\eta \int_Y k(\theta, y) \log \left(\frac{\mu_n k(y)}{p(y)} \right) \nu(\mathrm{d}y) \right]$$

→ NB : η corresponds to the learning rate

- Power descent : $\eta \in (0, 1]$, $(\alpha - 1)\kappa \geq 0$ and $\alpha \neq 1$

$$\Gamma(v) = [(\alpha - 1)v + 1]^{\frac{\eta}{1-\alpha}}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \left[\int_Y k(\theta, y) \left(\frac{\mu_n k(y)}{p(y)} \right)^{\alpha-1} \nu(\mathrm{d}y) + (\alpha - 1)\kappa \right]^{\frac{\eta}{1-\alpha}}$$

Examples satisfying (A2)

(A2) The function $\Gamma : \text{Dom}_\alpha \rightarrow \mathbb{R}_{>0}$ is decreasing, continuously differentiable and satisfies the inequality

$$[(\alpha - 1)(v - \kappa) + 1] (\log \Gamma)'(v) + 1 \geq 0 .$$

- Entropic Mirror Descent : $\eta \in (0, 1]$, $\kappa \in \mathbb{R}$ and $\alpha = 1$

$$\Gamma(v) = e^{-\eta v}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \exp \left[-\eta \int_Y k(\theta, y) \log \left(\frac{\mu_n k(y)}{p(y)} \right) \nu(\mathrm{d}y) \right]$$

→ NB : η corresponds to the learning rate

- Power descent : $\eta \in (0, 1]$, $(\alpha - 1)\kappa \geq 0$ and $\alpha \neq 1$

$$\Gamma(v) = [(\alpha - 1)v + 1]^{\frac{\eta}{1-\alpha}}$$

$$\mu_{n+1}(\mathrm{d}\theta) \propto \mu_n(\mathrm{d}\theta) \left[\int_Y k(\theta, y) \left(\frac{\mu_n k(y)}{p(y)} \right)^{\alpha-1} \nu(\mathrm{d}y) + (\alpha - 1)\kappa \right]^{\frac{\eta}{1-\alpha}}$$

Convergence results

Algorithm	Convergence results
<i>Entropic Mirror Descent</i> $\eta \in (0, \frac{1}{ \alpha-1 b _{\infty,\alpha}+1}), \kappa \in \mathbb{R}$	
<i>Power Descent</i> $\eta \in (0, 1], (\alpha - 1)\kappa \geqslant 0$	

Convergence results

Algorithm	Convergence results
<i>Entropic Mirror Descent</i> $\eta \in (0, \frac{1}{ \alpha-1 b _{\infty,\alpha}+1}), \kappa \in \mathbb{R}$	$O(1/N)$ convergence rates
<i>Power Descent</i> $\eta \in (0, 1], (\alpha - 1)\kappa \geqslant 0$	

Convergence results

Algorithm	Convergence results
<i>Entropic Mirror Descent</i> $\eta \in (0, \frac{1}{ \alpha-1 b _{\infty,\alpha}+1}), \kappa \in \mathbb{R}$	$O(1/N)$ convergence rates
<i>Power Descent</i> $\eta \in (0, 1], (\alpha - 1)\kappa \geqslant 0$	$\alpha > 1 : O(1/N)$ convergence rates

Convergence results

Algorithm	Convergence results
<i>Entropic Mirror Descent</i> $\eta \in (0, \frac{1}{ \alpha-1 b _{\infty,\alpha}+1}), \kappa \in \mathbb{R}$	$O(1/N)$ convergence rates
<i>Power Descent</i> $\eta \in (0, 1], (\alpha - 1)\kappa \geqslant 0$	$\alpha > 1 : O(1/N)$ convergence rates $\alpha < 1 : \text{convergence toward the optimum}$

The special case of finite mixture models

$$S_J = \left\{ \boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_J) \in \mathbb{R}^J : \forall j \in \{1, \dots, J\}, \lambda_j \geq 0 \text{ and } \sum_{j=1}^J \lambda_j = 1 \right\}$$

Let $\theta_1, \dots, \theta_J \in \mathcal{T}$ be **fixed** and denote

$$\mu_{\boldsymbol{\lambda}} = \sum_{j=1}^J \lambda_j \delta_{\theta_j} \quad \text{where } \boldsymbol{\lambda} \in S_J .$$

Then, $\mu_n = \underbrace{\mathcal{I}_{\alpha} \circ \cdots \circ \mathcal{I}_{\alpha}}_{n \text{ times}}(\mu_{\boldsymbol{\lambda}})$ is of the form $\mu_n = \sum_{j=1}^J \lambda_{j,n} \delta_{\theta_j}$ with

$$\begin{cases} \lambda_1 = \lambda \\ \lambda_{j,n+1} = \frac{\lambda_{j,n} \Gamma(b_{\mu_n, \alpha}(\theta_j) + \kappa)}{\sum_{i=1}^J \lambda_{i,n} \Gamma(b_{\mu_n, \alpha}(\theta_i) + \kappa)} . \end{cases}$$

→ In practice, we use Monte Carlo approximations to estimate $b_{\mu_n, \alpha}(\theta_j)$, e.g.

$$\hat{b}_{\mu_n, \alpha, M}(\theta_j) = \frac{1}{M} \sum_{m=1}^M \frac{k(\theta_j, Y_{m,n})}{\mu_n k(Y_{m,n})} f'_{\alpha} \left(\frac{\mu_n k(Y_{m,n})}{p(Y_{m,n})} \right),$$

with $Y_{1,n}, \dots, Y_{M,n} \stackrel{\text{i.i.d.}}{\sim} \mu_n k$.

→ NB : **Exploitation step** that does not require any information on the distribution of $\{\theta_1, \dots, \theta_J\}$

The special case of finite mixture models

$$S_J = \left\{ \boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_J) \in \mathbb{R}^J : \forall j \in \{1, \dots, J\}, \lambda_j \geq 0 \text{ and } \sum_{j=1}^J \lambda_j = 1 \right\}$$

Let $\theta_1, \dots, \theta_J \in \mathcal{T}$ be **fixed** and denote

$$\mu_{\boldsymbol{\lambda}} = \sum_{j=1}^J \lambda_j \delta_{\theta_j} \quad \text{where} \quad \boldsymbol{\lambda} \in S_J .$$

Then, $\mu_n = \underbrace{\mathcal{I}_{\alpha} \circ \cdots \circ \mathcal{I}_{\alpha}}_{n \text{ times}}(\mu_{\boldsymbol{\lambda}})$ is of the form $\mu_n = \sum_{j=1}^J \lambda_{j,n} \delta_{\theta_j}$ with

$$\begin{cases} \lambda_1 = \lambda \\ \lambda_{j,n+1} = \frac{\lambda_{j,n} \Gamma(b_{\mu_n, \alpha}(\theta_j) + \kappa)}{\sum_{i=1}^J \lambda_{i,n} \Gamma(b_{\mu_n, \alpha}(\theta_i) + \kappa)} . \end{cases}$$

→ In practice, we use Monte Carlo approximations to estimate $b_{\mu_n, \alpha}(\theta_j)$, e.g.

$$\hat{b}_{\mu_n, \alpha, M}(\theta_j) = \frac{1}{M} \sum_{m=1}^M \frac{k(\theta_j, Y_{m,n})}{\mu_n k(Y_{m,n})} f'_{\alpha} \left(\frac{\mu_n k(Y_{m,n})}{p(Y_{m,n})} \right),$$

with $Y_{1,n}, \dots, Y_{M,n} \stackrel{\text{i.i.d.}}{\sim} \mu_n k$.

→ NB : **Exploitation step** that does not require any information on the distribution of $\{\theta_1, \dots, \theta_J\}$

The special case of finite mixture models

$$S_J = \left\{ \boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_J) \in \mathbb{R}^J : \forall j \in \{1, \dots, J\}, \lambda_j \geq 0 \text{ and } \sum_{j=1}^J \lambda_j = 1 \right\}$$

Let $\theta_1, \dots, \theta_J \in \mathcal{T}$ be **fixed** and denote

$$\mu_{\boldsymbol{\lambda}} = \sum_{j=1}^J \lambda_j \delta_{\theta_j} \quad \text{where} \quad \boldsymbol{\lambda} \in S_J .$$

Then, $\mu_n = \underbrace{\mathcal{I}_{\alpha} \circ \cdots \circ \mathcal{I}_{\alpha}}_{n \text{ times}}(\mu_{\boldsymbol{\lambda}})$ is of the form $\mu_n = \sum_{j=1}^J \lambda_{j,n} \delta_{\theta_j}$ with

$$\begin{cases} \lambda_1 = \lambda \\ \lambda_{j,n+1} = \frac{\lambda_{j,n} \Gamma(b_{\mu_n, \alpha}(\theta_j) + \kappa)}{\sum_{i=1}^J \lambda_{i,n} \Gamma(b_{\mu_n, \alpha}(\theta_i) + \kappa)} . \end{cases}$$

→ In practice, we use Monte Carlo approximations to estimate $b_{\mu_n, \alpha}(\theta_j)$, e.g.

$$\hat{b}_{\mu_n, \alpha, M}(\theta_j) = \frac{1}{M} \sum_{m=1}^M \frac{k(\theta_j, Y_{m,n})}{\mu_n k(Y_{m,n})} f'_{\alpha} \left(\frac{\mu_n k(Y_{m,n})}{p(Y_{m,n})} \right),$$

with $Y_{1,n}, \dots, Y_{M,n} \stackrel{\text{i.i.d.}}{\sim} \mu_n k$.

→ NB : **Exploitation step** that does not require any information on the distribution of $\{\theta_1, \dots, \theta_J\}$

The special case of finite mixture models

$$S_J = \left\{ \boldsymbol{\lambda} = (\lambda_1, \dots, \lambda_J) \in \mathbb{R}^J : \forall j \in \{1, \dots, J\}, \lambda_j \geq 0 \text{ and } \sum_{j=1}^J \lambda_j = 1 \right\}$$

Let $\theta_1, \dots, \theta_J \in \mathcal{T}$ be **fixed** and denote

$$\mu_{\boldsymbol{\lambda}} = \sum_{j=1}^J \lambda_j \delta_{\theta_j} \quad \text{where} \quad \boldsymbol{\lambda} \in S_J .$$

Then, $\mu_n = \underbrace{\mathcal{I}_{\alpha} \circ \dots \circ \mathcal{I}_{\alpha}}_{n \text{ times}}(\mu_{\boldsymbol{\lambda}})$ is of the form $\mu_n = \sum_{j=1}^J \lambda_{j,n} \delta_{\theta_j}$ with

$$\begin{cases} \lambda_1 = \lambda \\ \lambda_{j,n+1} = \frac{\lambda_{j,n} \Gamma(b_{\mu_n, \alpha}(\theta_j) + \kappa)}{\sum_{i=1}^J \lambda_{i,n} \Gamma(b_{\mu_n, \alpha}(\theta_i) + \kappa)} . \end{cases}$$

→ In practice, we use Monte Carlo approximations to estimate $b_{\mu_n, \alpha}(\theta_j)$, e.g.

$$\hat{b}_{\mu_n, \alpha, M}(\theta_j) = \frac{1}{M} \sum_{m=1}^M \frac{k(\theta_j, Y_{m,n})}{\mu_n k(Y_{m,n})} f'_{\alpha} \left(\frac{\mu_n k(Y_{m,n})}{p(Y_{m,n})} \right),$$

with $Y_{1,n}, \dots, Y_{M,n} \stackrel{\text{i.i.d.}}{\sim} \mu_n k$.

→ NB : **Exploitation step** that does not require any information on the distribution of $\{\theta_1, \dots, \theta_J\}$

Outline

- 1 Introduction
- 2 Infinite-dimensional α -divergence minimisation
- 3 Numerical experiments
- 4 Conclusion

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbb{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbb{T}^J \right\}.$$

Full algorithm

- ➊ Exploitation step : optimise λ using the (α, Γ) -descent.
- ➋ Exploration step : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(\mathbf{y}) = Z \times [0.5\mathcal{N}(\mathbf{y}; -2\mathbf{u}_d, \mathbf{I}_d) + 0.5\mathcal{N}(\mathbf{y}; 2\mathbf{u}_d, \mathbf{I}_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbf{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbf{T}^J \right\}.$$

Full algorithm

- ➊ Exploitation step : optimise λ using the (α, Γ) -descent.
- ➋ Exploration step : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(\mathbf{y}) = Z \times [0.5\mathcal{N}(\mathbf{y}; -2\mathbf{u}_d, \mathbf{I}_d) + 0.5\mathcal{N}(\mathbf{y}; 2\mathbf{u}_d, \mathbf{I}_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbb{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbb{T}^J \right\}.$$

Full algorithm

① **Exploitation step** : optimise λ using the (α, Γ) -descent.

② Exploration step : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(\mathbf{y}) = Z \times [0.5\mathcal{N}(\mathbf{y}; -2\mathbf{u}_d, \mathbf{I}_d) + 0.5\mathcal{N}(\mathbf{y}; 2\mathbf{u}_d, \mathbf{I}_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbb{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbb{T}^J \right\}.$$

Full algorithm

- ① **Exploitation step** : optimise λ using the (α, Γ) -descent.
- ② **Exploration step** : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(y) = Z \times [0.5\mathcal{N}(y; -2u_d, I_d) + 0.5\mathcal{N}(y; 2u_d, I_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbb{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbb{T}^J \right\}.$$

Full algorithm

- ➊ **Exploitation step** : optimise λ using the (α, Γ) -descent.
- ➋ **Exploration step** : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(\mathbf{y}) = Z \times [0.5\mathcal{N}(\mathbf{y}; -2\mathbf{u}_d, \mathbf{I}_d) + 0.5\mathcal{N}(\mathbf{y}; 2\mathbf{u}_d, \mathbf{I}_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Numerical experiments

- Gaussian kernel with density k_h and bandwidth h , $\mathbb{T} = \mathbb{R}^d$

$$\left\{ y \mapsto \mu_{\lambda, \Theta} k_h(y) = \sum_{j=1}^J \lambda_j k_h(y - \theta_j) : \lambda \in \mathcal{S}_J, \Theta \in \mathbb{T}^J \right\}.$$

Full algorithm

- ➊ Exploitation step : optimise λ using the (α, Γ) -descent.
- ➋ Exploration step : update Θ (e.g. by sampling under $\mu_{\lambda, \Theta} k_h$, $h \propto J^{-1/(4+d)}$)

- Toy example

$$p(\mathbf{y}) = Z \times [0.5\mathcal{N}(\mathbf{y}; -2\mathbf{u}_d, \mathbf{I}_d) + 0.5\mathcal{N}(\mathbf{y}; 2\mathbf{u}_d, \mathbf{I}_d)], Z = 2$$

- Bayesian Logistic Regression Covertype dataset (581,012 data points and 54 features)

Toy example : Entropic Mirror Descent vs Power Descent

Comparison between

- 0.5-Mirror descent : $\Gamma(v) = e^{-\eta v}$ and $\alpha = 0.5$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ and $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$


$\eta_n = \eta_0 / \sqrt{n}$, $\eta_0 = 0.5$, cv criterion : VR-Bound averaged over 100 trials

Toy example : Entropic Mirror Descent vs Power Descent

Comparison between

- 0.5-Mirror descent : $\Gamma(v) = e^{-\eta v}$ and $\alpha = 0.5$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ and $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0 / \sqrt{n}$, $\eta_0 = 0.5$, cv criterion : VR-Bound averaged over 100 trials




Toy example : Entropic Mirror Descent vs Power Descent

Comparison between

- 0.5-Mirror descent : $\Gamma(v) = e^{-\eta v}$ and $\alpha = 0.5$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ and $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0/\sqrt{n}$, $\eta_0 = 0.5$, cv criterion : VR-Bound averaged over 100 trials






Toy example : Entropic Mirror Descent vs Power Descent

Comparison between

- 0.5-Mirror descent : $\Gamma(v) = e^{-\eta v}$ and $\alpha = 0.5$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ and $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0/\sqrt{n}$, $\eta_0 = 0.5$, cv criterion : VR-Bound averaged over 100 trials



Toy example : the case $\alpha = 1$

Comparison between:

- 1-Mirror descent : $\Gamma(v) = e^{-\eta v}$ with $\alpha = 1$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ with $\alpha = 0.5$.


$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0 / \sqrt{n}$, $\eta_0 = 0.5$, cv criterion : llh averaged over 100 trials

Toy example : the case $\alpha = 1$

Comparison between:

- 1-Mirror descent : $\Gamma(v) = e^{-\eta v}$ with $\alpha = 1$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ with $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0/\sqrt{n}$, $\eta_0 = 0.5$, cv criterion : llh averaged over 100 trials




Toy example : the case $\alpha = 1$

Comparison between:

- 1-Mirror descent : $\Gamma(v) = e^{-\eta v}$ with $\alpha = 1$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ with $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0/\sqrt{n}$, $\eta_0 = 0.5$, cv criterion : llh averaged over 100 trials






Toy example : the case $\alpha = 1$

Comparison between:

- 1-Mirror descent : $\Gamma(v) = e^{-\eta v}$ with $\alpha = 1$,
- 0.5-Power descent : $\Gamma(v) = [(\alpha - 1)v + 1]^{\eta/(1-\alpha)}$ with $\alpha = 0.5$.

$J = M = 100$, initial mixture weights : $[1/J, \dots, 1/J]$, $N = 10$, $T = 20$
 $\eta_n = \eta_0/\sqrt{n}$, $\eta_0 = 0.5$, cv criterion : llh averaged over 100 trials



Bayesian Logistic Regression

→ $\mathcal{D} = \{\mathbf{c}, \mathbf{x}\}$: I binary class labels, $c_i \in \{-1, 1\}$, L covariates for each datapoint, $\mathbf{x}_i \in \mathbb{R}^L$

→ Model : L regression coefficients $w_l \in \mathbb{R}$, precision parameter $\beta \in \mathbb{R}^+$

$$p_0(\beta) = \text{Gamma}(\beta; a, b),$$

$$p_0(w_l | \beta) = \mathcal{N}(w_l; 0, \beta^{-1}), \quad 1 \leq l \leq L$$

$$p(c_i = 1 | \mathbf{x}_i, \mathbf{w}) = \frac{1}{1 + e^{-\mathbf{w}^T \mathbf{x}_i}}, \quad 1 \leq i \leq I$$

where $a = 1$ and $b = 0.01$

Nonparametric variational inference S. Gershman, M. Hoffman, and D. Blei (2012). ICML

→ Quantity of interest : $p(y|\mathcal{D})$ with $y = [\mathbf{w}, \log \beta]$

Comparison between

- 0.5-Power descent
- Typical AIS

$$N = 1, T = 500, J_0 = M_0 = 20, J_{t+1} = M_{t+1} = J_t + 1$$

initial mixture weights : $[1/J_t, \dots, 1/J_t]$, $\eta_n = \eta_0/\sqrt{n}$ with $\eta_0 = 0.05$

Bayesian Logistic Regression

→ $\mathcal{D} = \{\mathbf{c}, \mathbf{x}\}$: I binary class labels, $c_i \in \{-1, 1\}$, L covariates for each datapoint, $\mathbf{x}_i \in \mathbb{R}^L$

→ Model : L regression coefficients $w_l \in \mathbb{R}$, precision parameter $\beta \in \mathbb{R}^+$

$$p_0(\beta) = \text{Gamma}(\beta; a, b),$$

$$p_0(w_l | \beta) = \mathcal{N}(w_l; 0, \beta^{-1}), \quad 1 \leq l \leq L$$

$$p(c_i = 1 | \mathbf{x}_i, \mathbf{w}) = \frac{1}{1 + e^{-\mathbf{w}^T \mathbf{x}_i}}, \quad 1 \leq i \leq I$$

where $a = 1$ and $b = 0.01$

Nonparametric variational inference S. Gershman, M. Hoffman, and D. Blei (2012). ICML

→ Quantity of interest : $p(y|\mathcal{D})$ with $y = [\mathbf{w}, \log \beta]$

Comparison between

- 0.5-Power descent
- Typical AIS

$$N = 1, T = 500, J_0 = M_0 = 20, J_{t+1} = M_{t+1} = J_t + 1$$

initial mixture weights : $[1/J_t, \dots, 1/J_t]$, $\eta_n = \eta_0/\sqrt{n}$ with $\eta_0 = 0.05$

Bayesian Logistic Regression

→ $\mathcal{D} = \{\mathbf{c}, \mathbf{x}\}$: I binary class labels, $c_i \in \{-1, 1\}$, L covariates for each datapoint, $\mathbf{x}_i \in \mathbb{R}^L$

→ Model : L regression coefficients $w_l \in \mathbb{R}$, precision parameter $\beta \in \mathbb{R}^+$

$$p_0(\beta) = \text{Gamma}(\beta; a, b),$$

$$p_0(w_l | \beta) = \mathcal{N}(w_l; 0, \beta^{-1}), \quad 1 \leq l \leq L$$

$$p(c_i = 1 | \mathbf{x}_i, \mathbf{w}) = \frac{1}{1 + e^{-\mathbf{w}^T \mathbf{x}_i}}, \quad 1 \leq i \leq I$$

where $a = 1$ and $b = 0.01$

Nonparametric variational inference S. Gershman, M. Hoffman, and D. Blei (2012). ICML

→ Quantity of interest : $p(y|\mathcal{D})$ with $y = [\mathbf{w}, \log \beta]$

Comparison between

- 0.5-Power descent
- Typical AIS

$$N = 1, T = 500, J_0 = M_0 = 20, J_{t+1} = M_{t+1} = J_t + 1$$

initial mixture weights : $[1/J_t, \dots, 1/J_t]$, $\eta_n = \eta_0/\sqrt{n}$ with $\eta_0 = 0.05$

Bayesian Logistic Regression

→ $\mathcal{D} = \{\mathbf{c}, \mathbf{x}\}$: I binary class labels, $c_i \in \{-1, 1\}$, L covariates for each datapoint, $\mathbf{x}_i \in \mathbb{R}^L$

→ Model : L regression coefficients $w_l \in \mathbb{R}$, precision parameter $\beta \in \mathbb{R}^+$

$$p_0(\beta) = \text{Gamma}(\beta; a, b),$$

$$p_0(w_l | \beta) = \mathcal{N}(w_l; 0, \beta^{-1}), \quad 1 \leq l \leq L$$

$$p(c_i = 1 | \mathbf{x}_i, \mathbf{w}) = \frac{1}{1 + e^{-\mathbf{w}^T \mathbf{x}_i}}, \quad 1 \leq i \leq I$$



where $a = 1$ and $b = 0.01$

Nonparametric variational inference S. Gershman, M. Hoffman, and D. Blei (2012). ICML

→ Quantity of interest : $p(y|\mathcal{D})$ with $y = [\mathbf{w}, \log \beta]$

Comparison between

- 0.5-Power descent
- Typical AIS



$N = 1, T = 500, J_0 = M_0 = 20, J_{t+1} = M_{t+1} = J_t + 1$

initial mixture weights : $[1/J_t, \dots, 1/J_t]$, $\eta_n = \eta_0/\sqrt{n}$ with $\eta_0 = 0.05$

Outline

- 1 Introduction
- 2 Infinite-dimensional α -divergence minimisation
- 3 Numerical experiments
- 4 Conclusion

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the Entropic Mirror Descent algorithm
- novel Power Descent algorithm
- conditions for a systematic decrease + convergence results
- applicable to mixture models :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ not specified (e.g. your favorite update for Θ)
- ② Empirical advantages of using the Power Descent algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the **Entropic Mirror Descent** algorithm
- novel **Power Descent** algorithm
- conditions for a systematic decrease + convergence results
- applicable to mixture models :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ not specified (e.g. your favorite update for Θ)
- ② Empirical advantages of using the **Power Descent** algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the **Entropic Mirror Descent** algorithm
- novel **Power Descent** algorithm
- conditions for a systematic decrease + convergence results
- applicable to mixture models :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ not specified (e.g. your favorite update for Θ)
- ② Empirical advantages of using the **Power Descent** algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the **Entropic Mirror Descent** algorithm
- novel **Power Descent** algorithm
- conditions for a **systematic decrease + convergence results**
- applicable to mixture models :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ not specified (e.g. your favorite update for Θ)
- ② Empirical advantages of using the **Power Descent** algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathsf{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathsf{M} \right\}$$

- recovers the [Entropic Mirror Descent](#) algorithm
- novel [Power Descent](#) algorithm
- conditions for a [systematic decrease](#) + [convergence](#) results
- applicable to [mixture models](#) :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathsf{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ [not specified](#) (e.g. [your](#) favorite update for Θ)
- ② Empirical advantages of using the [Power Descent](#) algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathsf{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathsf{M} \right\}$$

- recovers the [Entropic Mirror Descent](#) algorithm
- novel [Power Descent](#) algorithm
- conditions for a [systematic decrease](#) + [convergence](#) results
- applicable to [mixture models](#) :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathsf{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ [not specified](#) (e.g. [your](#) favorite update for Θ)
- ② Empirical advantages of using the [Power Descent](#) algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the **Entropic Mirror Descent** algorithm
- novel **Power Descent** algorithm
- conditions for a **systematic decrease** + **convergence** results
- applicable to **mixture models** :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ➊ Update for Θ **not specified** (e.g. **your** favorite update for Θ)
- ➋ Empirical advantages of using the **Power Descent** algorithm

Summary

General framework for infinite-dimensional α -divergence minimisation over

$$\mathcal{Q} = \left\{ q : y \mapsto \int_{\mathcal{T}} \mu(d\theta) k(\theta, y) : \mu \in \mathcal{M} \right\}$$

- recovers the **Entropic Mirror Descent** algorithm
- novel **Power Descent** algorithm
- conditions for a **systematic decrease** + **convergence** results
- applicable to **mixture models** :

$$\mathcal{Q} = \left\{ q : y \mapsto \sum_{j=1}^J \lambda_j k(\theta_j, y) : \boldsymbol{\lambda} \in \mathcal{S}_J, \Theta \in \mathcal{T}^J \right\}$$

→ Exploitation - Exploration algorithm

- ① Update for Θ **not specified** (e.g. **your** favorite update for Θ)
- ② Empirical advantages of using the **Power Descent** algorithm

Extensions

① Mixture weights optimisation for Alpha-Divergence Variational Inference.

K. Daudel and R. Douc (2021). To appear in NeurIPS2021

→ Extension of the Power Descent to the case $\alpha = 1$

→ Full proof of convergence for finite mixture models ($\alpha < 1$)

→ Closely-related algorithm : Rényi Descent

② Monotonic Alpha-divergence Minimisation.

K. Daudel, R. Douc and F. Roueff (2021). <https://arxiv.org/abs/2103.05684>

→ Conditions for a simultaneous optimisation w.r.t λ and Θ
(that preserve the monotonic decrease!)

→ Simple updates on Θ for specific kernels k (e.g Gaussian, Student's)

→ Links with Gradient Descent schemes and an Integrated EM algorithm
with empirical benefits

③ Your next paper?

Extensions

① Mixture weights optimisation for Alpha-Divergence Variational Inference.

K. Daudel and R. Douc (2021). To appear in NeurIPS2021

→ Extension of the Power Descent to the case $\alpha = 1$

→ Full proof of convergence for finite mixture models ($\alpha < 1$)

→ Closely-related algorithm : Rényi Descent

② Monotonic Alpha-divergence Minimisation.

K. Daudel, R. Douc and F. Roueff (2021). <https://arxiv.org/abs/2103.05684>

→ Conditions for a simultaneous optimisation w.r.t λ and Θ
(that preserve the monotonic decrease!)

→ Simple updates on Θ for specific kernels k (e.g Gaussian, Student's)

→ Links with Gradient Descent schemes and an Integrated EM algorithm
with empirical benefits

③ Your next paper?

Extensions

① Mixture weights optimisation for Alpha-Divergence Variational Inference.

K. Daudel and R. Douc (2021). To appear in NeurIPS2021

→ Extension of the Power Descent to the case $\alpha = 1$

→ Full proof of convergence for finite mixture models ($\alpha < 1$)

→ Closely-related algorithm : Rényi Descent

② Monotonic Alpha-divergence Minimisation.

K. Daudel, R. Douc and F. Roueff (2021). <https://arxiv.org/abs/2103.05684>

→ Conditions for a simultaneous optimisation w.r.t λ and Θ
(that preserve the monotonic decrease!)

→ Simple updates on Θ for specific kernels k (e.g Gaussian, Student's)

→ Links with Gradient Descent schemes and an Integrated EM algorithm
with empirical benefits

③ Your next paper?

Extensions

① Mixture weights optimisation for Alpha-Divergence Variational Inference.

K. Daudel and R. Douc (2021). To appear in NeurIPS2021

→ Extension of the Power Descent to the case $\alpha = 1$

→ Full proof of convergence for finite mixture models ($\alpha < 1$)

→ Closely-related algorithm : Rényi Descent

② Monotonic Alpha-divergence Minimisation.

K. Daudel, R. Douc and F. Roueff (2021). <https://arxiv.org/abs/2103.05684>

→ Conditions for a simultaneous optimisation w.r.t λ and Θ
(that preserve the monotonic decrease!)

→ Simple updates on Θ for specific kernels k (e.g Gaussian, Student's)

→ Links with Gradient Descent schemes and an Integrated EM algorithm
with empirical benefits

③ Your next paper?

Extensions

① Mixture weights optimisation for Alpha-Divergence Variational Inference.

K. Daudel and R. Douc (2021). To appear in NeurIPS2021

→ Extension of the Power Descent to the case $\alpha = 1$

→ Full proof of convergence for finite mixture models ($\alpha < 1$)

→ Closely-related algorithm : Rényi Descent

② Monotonic Alpha-divergence Minimisation.

K. Daudel, R. Douc and F. Roueff (2021). <https://arxiv.org/abs/2103.05684>

→ Conditions for a simultaneous optimisation w.r.t λ and Θ
(that preserve the monotonic decrease!)

→ Simple updates on Θ for specific kernels k (e.g Gaussian, Student's)

→ Links with Gradient Descent schemes and an Integrated EM algorithm
with empirical benefits

③ Your next paper?

Thank you for your attention!

kamelia.daudel@stats.ox.ac.uk